

forcing occurs in C^n until the cut-vertex v is turned black. Now, if v is the last vertex on C^{n-1} to be turned black, then it's clear that at most two vertices in $\langle V(C^{n-1} \cup C^n) \rangle$ are turned black at a time – since, obviously, with any $Z(B_n)$ -set at most two vertices are turned black on any cycle C^i at each step. On the other hand, if v is turned black before C^{n-1} is turned entirely black, then v , being a neighbor to at least two white vertices, can not force until C^{n-1} is turned entirely black. We've thus shown that $Z_m \cap V(C^{n-1} \cup C^n)$ contains at most two forcing vertices for any m or, equivalently, at most two vertices in $\langle V(C^{n-1} \cup C^n) \rangle$ are turned black at a time. \square

Acknowledgement. The authors wish to thank the anonymous referee for some suggestions and corrections.

References

- [1] AIM Minimum Rank - Special Graphs Work Group (F. Barioli, W. Barrett, S. Butler, S. M. Cioabă, D. Cvetković, S. M. Fallat, C. Godsil, W. Haemers, L. Hogben, R. Mikkelsen, S. Narayan, O. Pryporova, I. Sciriha, W. So, D. Stevanović, H. van der Holst, K. Vander Meulen, A. Wangsness). Zero forcing sets and the minimum rank of graphs. *Linear Algebra and its Applications*, **428/7** (2008) 1628-1648.
- [2] F. Barioli, W. Barrett, S. M. Fallat, H. T. Hall, L. Hogben, B. Shader, P. van den Driessche, and H. van der Holst, Zero forcing parameters and minimum rank problems, *preprint* (2010).
- [3] A. Berman, S. Friedland, L. Hogben, U. G. Rothblum, and B. Shader, An upper bound for the minimum rank of a graph. *Linear Algebra Appl.*, **429** (2008) 1629-1638.
- [4] G. Chartrand and P. Zhang, *Introduction to Graph Theory*. McGraw-Hill, Kalamazoo, MI (2004).
- [5] C. J. Edholm, L. Hogben, M. Hyunh, J. LaGrange, and D. D. Row, Vertex and edge spread of zero forcing number, maximum nullity, and minimum rank of a graph. *preprint* (2010).
- [6] L. Huang, G. J. Chang, and H. Yeh, On minimum rank and zero forcing sets of a graph. *Linear Algebra and its Applications*, **432** (2010) 2961-2973.
- [7] C. R. Johnson and A. Leal Duarte, The maximum multiplicity of an eigenvalue in a matrix whose graph is a tree. *Linear Multilinear Algebra*, **46** (1999) 139-144.
- [8] J. Llibre and M. Todd, Periods, Lefschetz numbers and entropy for a class of maps on a bouquet of circles, arXiv:math/0409361v1 [math.DS]
- [9] W. S. Massey, *Algebraic Topology: An Introduction*. Graduate Texts in Mathematics **56**, Springer-Verlag (1989)

REMARKS ON THE DOMINATION NUMBER OF GRAPHS

CSABA BIRÓ, ÉVA CZABARKA, PETER DANKELMANN,
AND LÁSZLÓ SZÉKELY

ABSTRACT. This paper consists of two loosely related notes on the domination number of graphs. In the first part, we provide a new upper bound for the domination number of d -regular graphs. Our bound is the best known for $d \geq 6$. In the second part, we compute the domatic number of certain Kneser graphs.

1. INTRODUCTION

Let $G = (V, E)$ be a graph and $X \subseteq V$. X is a dominating set if for every $v \in V$ the closed neighborhood of v and X are not disjoint. X is a total dominating set if for every $v \in V$ the open neighborhood of v and X are not disjoint.

The domination number $\gamma(G)$ is the minimum size of a dominating set and the total domination number $\gamma_t(G)$ is the minimum size of a total dominating set. It is clear from the definition that $\gamma(G) \leq \gamma_t(G) \leq 2\gamma(G)$.

A domatic partition of a graph G is a partition of V into sets V_1, \dots, V_k such that each V_i is a dominating set of G . The domatic number $\text{dom}(G)$ of a graph G is the maximum size of a domatic partition. From the definition it is immediate that $\text{dom}(G) \leq \frac{|V(G)|}{\gamma(G)}$.

Extensive research has been done to find upper bounds for the domination number of graphs of given minimum degree. Alon and Spencer [2] provide a probabilistic proof for the following theorem.

Theorem 1.1. *Let G be a graph on n vertices with minimum degree $\delta > 1$. Then*

$$(1) \quad \gamma(G) \leq \frac{1 + \ln(\delta + 1)}{\delta + 1} \cdot n$$

The second authors was partially supported by a grant from the University of South Carolina Promising Investigator Research Award and enjoyed the hospitality of the Rheinische Friedrich-Wilhelms Universität, Bonn. The fourth author was supported in part by the NSF DMS contracts 7001111 and 1000645; and the Alexander von Humboldt Stiftung at the Rheinische Friedrich-Wilhelms Universität, Bonn.

Arnautov [3], Lovász [12], [13], and Payan [17] obtained a slightly better bound by considering a dominating set constructed by the following greedy algorithm: select the dominating set vertex by vertex in such a way that in every step we choose a vertex that dominates the maximum number of yet undominated vertices. (If there are more than one choices, we choose randomly.) They obtained

$$(2) \quad \gamma(G) \leq \frac{n}{\delta+1} H_{\delta+1},$$

where $H_k = 1 + \frac{1}{2} + \dots + \frac{1}{k}$ is the k^{th} harmonic number.

It can be deduced from the results in Alon [1] that these bounds are asymptotically optimal as $\delta \rightarrow \infty$. Nevertheless, the question of the optimal bound for the domination number is still interesting for small minimum degrees.

Another, related question is to study the optimal bound for regular graphs. It is obvious that for a d -regular graph G on n vertices we have

$$(3) \quad \gamma(G) \geq \frac{1}{d+1} \cdot n.$$

Denoting the class of d -regular graphs by \mathcal{G}_d , we may define the parameter $D(d)$ as

$$(4) \quad D(d) = \sup \left\{ \frac{\gamma(G)}{|V(G)|} : G \in \mathcal{G}_d \right\} = \inf \{t : G \in \mathcal{G}_d, \gamma(G) \leq t|V(G)|\}.$$

As $D(d)$ gives an upper bound for the dominating number of d -regular graphs, it is a natural problem to find good lower/upper bounds for $D(d)$. We list the best available information about $D(d)$ for small d in Table 1 with references to the first publication of the relevant results.

For comparison, we note the best known upper bounds for the domination number of graphs of given small fixed minimum degree. Let G be a graph of order n . Ore [16] showed that if $\delta(G) \geq 1$, then $\gamma(G) \leq n/2$. McCuaig and Shepherd [14] showed that if $\delta(G) \geq 2$ (and G is not one of seven exceptional graphs), then $\gamma(G) \leq (2/5)n$. Reed [18] showed that if $\delta(G) \geq 3$, then $\gamma(G) \leq (3/8)n$. In the monograph [9] by Haynes, Hedetniemi and Slater there appeared the general conjecture that for $\delta(G) = k \geq 4$, $\gamma(G) \leq kn/(3k-1)$. The conjecture was proven in a sequence of papers, and in fact, for large values k , stronger inequalities are satisfied.

In Section 2 we prove a new upper bound for the dominating number for d -regular graphs. Our bound is significantly stronger than (1) in case of small $\delta = d$. While better bounds are known for $d < 6$ (see Table 1), our bound is the strongest known for any fixed $d \geq 6$.

In Section 3 we compute some exact values of the domatic number of certain Kneser graphs. Originally we hoped that odd graphs (i.e. Kneser graphs $K(2n+1, n)$, studied in [15]), which are $(n+1)$ -regular, may provide

d	$D(d)$
1	$\frac{1}{2}$
2	$\frac{1}{2}$
3	$\frac{3}{8}$ [18, 5]
4	$\leq \frac{4}{11}$ [11]
5	$\leq \frac{5}{14}$ [19]

TABLE 1. Upper bound for the domination number for small d

improved lower bounds for $D(d)$, however, somewhat surprisingly, this is not the case for small d due to the existence of certain block designs.

2. UPPER BOUND FOR γ FOR d -REGULAR GRAPHS

Clark et al. [6] showed that the greedy algorithm proposed by Alon and Spencer in [2] can be used to give better bounds on the domination number of graphs with a fixed minimum degree. They also found further improvement for regular graphs. They showed that if $V(G) = n$ for a d -regular graph G , then

$$(5) \quad \gamma(G) \leq C_d n \quad \text{where} \quad C_d = 1 - \frac{d-1 + \frac{1}{d^2+1}}{\prod_{j=1}^{d-1} (1 + \frac{d+1}{jd})}.$$

We prove a similar theorem with a strictly smaller constant. (Strictly speaking, Clark et al. [6] did not provide a proof to the improvement for d -regular graphs, just argued—correctly—that it is analogous to the proof in the minimum degree condition. If one accepts this claim, then he should our improvement as well, without a detailed proof. For the sake of completeness, however, we provide the proof.) Our main result is the following.

Theorem 2.1. For a d -regular graph G on n vertices,

$$(6) \quad \gamma(G) \leq B_d n \quad \text{where} \quad B_d = 1 - \frac{d^2 - d + 1}{1 + d \prod_{j=1}^{d-1} \left(1 + \frac{d+1}{jd}\right)}.$$

Just from the formulas, it is not obvious that $B_d < C_d$. However, the proof easily implies this fact, and it is also easy to compare the constants for specific values of d , as they are both rational numbers. E.g. $C_6 = \frac{570971}{1690715} \approx 0.3377 > B_6 = \frac{483425}{1447649} \approx 0.3339$.

The theorem works for all $d \geq 1$, but, as mentioned in the introduction, for $d < 6$ there are stronger or equally strong results in the literature, listed in Table 1. Asymptotic analysis of the bounds are shown in Table 2.

$$\begin{aligned}
& \frac{\ln d+1}{d} - O\left(\frac{\ln d}{d^2}\right) \\
& + \frac{\ln d+\gamma}{d} - \frac{\ln d-\frac{3}{2}+\gamma}{d^2} + \frac{12 \ln d+12\gamma-31}{12d^3} - O\left(\frac{\ln d}{d^4}\right) \\
& + \frac{\ln d+\gamma}{d} - \frac{\frac{1}{2}(\ln d)^2+\gamma \ln d-\frac{3}{2}+\frac{\pi^2+6\gamma^2}{12}}{d^2} + \frac{2(\ln d)^3+6\gamma(\ln d)^2+(6\gamma^2+\pi^2-18) \ln d+4\zeta(3)+\pi^2\gamma+2\gamma^3-7-18\gamma}{12d^3} + O\left(\frac{(\ln d)^4}{d^4}\right) \\
& + \frac{\ln d+\gamma}{d} - \frac{\frac{1}{2}(\ln d)^2+\gamma \ln d-\frac{3}{2}+\frac{\pi^2+6\gamma^2}{12}}{d^2} + \frac{2(\ln d)^3+6\gamma(\ln d)^2+(6\gamma^2+\pi^2-30) \ln d+4\zeta(3)+\pi^2\gamma+2\gamma^3+5-30\gamma}{12d^3} + O\left(\frac{(\ln d)^4}{d^4}\right)
\end{aligned}$$

TABLE 2. For comparison, we include here asymptotic formulae for (1) [Alon [2]], (2) [Arnautov [3], Lovász [12], [13] and Payan [17]], (5) [Clark et al. [6]], and our bound (6), in this order, on the fraction $D(d)$ as function of d , as $d \rightarrow \infty$. We obtained these results with Maple. The first two bounds hold for graphs with minimum degree d as well. The Euler-Mascheroni constant is denoted by $\gamma = 0.5772\dots$, the Riemann zeta function in 3 is $\zeta(3)$. The successive improvements always happen in the coefficient of the next power of d .

2.1. Proof of Theorem 2.1. Let G be a graph and $T \subseteq V(G)$. One can build a dominating set of G starting from T and then using the greedy algorithm described above to add vertices to T , until a dominating set is reached. The following lemma is essentially an adapted form of the main argument in [6], keeping the same notation as far as possible for easy comparison.

Lemma 2.1.1. *Let G be a graph of minimum degree δ and a maximal $T \subseteq V(G)$ such that any two vertices in T are of distance at least 3. Let $n = |V(G)|$ and $t = |T|/n$. Suppose that the greedy algorithm builds a dominating set D starting from T . Then*

$$|D| \leq A_\delta n$$

where

$$A_\delta = 1 - \frac{\delta - 1 + t}{\prod_{j=1}^{\delta-1} \left(1 + \frac{\delta+1}{j\delta}\right)}.$$

Proof. Suppose that after a certain step in the greedy algorithm, the number of yet uncovered vertices is r , and the size of the built set is i . Each of the yet uncovered vertices have a closed neighborhood of cardinality at least $\delta + 1$, so their total cardinalities is at least $r(\delta + 1)$. The $n - i$ points that are not in the built set are on average in $r(\delta + 1)/(n - i)$ of these neighborhoods, so there exists a vertex that has at least $r(\delta + 1)/(n - i)$ uncovered neighbor. Note that this vertex may be covered. The arguments shows that in every new step, we decrease the number of uncovered vertices by at least $r(\delta + 1)/(n - i)$.

Define $g_0 = n - (\delta + 1)T$. We define the sequence g_ℓ , which is an upper bound on the number of not yet dominated vertices after ℓ greedy choices to extend T into a dominating set. For $\ell > 0$, we know from the averaging argument above that if there are r not yet dominated vertices after ℓ greedy steps, then after a greedy choice to dominate as much as possible, at most $r \left(1 - \frac{\delta+1}{n-\ell-|T|}\right)$ are not dominated, for any $\ell > 0$. Hence we set $g_{\ell+1} = \left\lfloor g_\ell \left(1 - \frac{\delta+1}{n-\ell-|T|}\right) \right\rfloor$. Our analogue of the $\gamma_6(G)$ bound for the domination number of G in [6] is $\gamma'_6(G) = |T| + \min\{\ell' : g_{\ell'} = 0\}$.

Observe that $g_{n-\delta} = 0$ and $g_0 = n - (\delta + 1)T > g_1 > g_2 > \dots > g_{\gamma'_6(G)} = 0$. Define

$$R_{\ell+1} = \left\lceil g_\ell \left(1 - \frac{\delta + 1}{n - \ell - |T|}\right) \right\rceil$$

for $0 \leq \ell < \gamma'_6(G)$ and observe that in this range $g_\ell > g_{\ell+1} = g_\ell - R_{\ell+1}$, and hence $R_{\ell+1} \geq 1$. As T was maximal, we conclude that

$$\delta \geq R_1 \geq R_2 \geq \dots \geq R_{\gamma'_6(G)} \geq 1.$$

For $2 \leq j \leq \delta + 1$, set

$$n_j = |\{i : 1 \leq i \leq \gamma'_6(G) \text{ with } R_i = \delta + 2 - j\}|.$$

Clearly $\gamma'_6(G) = |T| + n_2 + \dots + n_{\delta+1}$. We are going to find a recursive formula for the n_i numbers.

For $1 \leq \ell \leq n_2$, we have $g_\ell = n - (\delta + 1)|T| - \ell\delta$, and for any $0 \leq n' \leq n_k$ and $\ell = n_2 + \dots + n_{k-1} + n'$, we have

$$g_\ell = n - (\delta + 1)|T| - \sum_{j=2}^{k-1} (\delta + 2 - j)n_j - n'(\delta + 2 - k).$$

It is clear that n_2 is the least integer such that $R_{n_2+1} \leq \delta - 1$. Using the definition of R_{n_2+1} and the displayed formula above for g_{n_2} , this means

$$\left(n - (\delta + 1)|T| - N_2\delta \right) \frac{\delta + 1}{n - n_2 - |T|} \leq \delta - 1,$$

which is equivalent to

$$\frac{2n - |T|(\delta^2 + \delta + 2)}{\delta^2 + 1} \leq n_2.$$

$$\text{Hence } n_2 = \left\lceil \frac{2n - |T|(\delta^2 + \delta + 2)}{\delta^2 + 1} \right\rceil.$$

With a similar argument, for $2 < k \leq \delta + 1$, we obtain that n_k is the least integer satisfying $R_{n_2+\dots+n_{k+1}} \leq \delta + 2 - (k + 1)$, which by a similar argument corresponds to

$$\left(n - (\delta + 1)|T| - \sum_{j=2}^k (\delta + 2 - j)n_j \right) \frac{\delta + 1}{n - |T| - (n_2 + \dots + n_k)} = g_{n_2+\dots+n_k} \frac{\delta + 1}{n - |T| - (n_2 + \dots + n_k)} \leq \delta + 2 - (k + 1).$$

Solving the inequality above for n_k , we obtain after some algebra

$$\frac{kn - (\delta^2 + \delta + 2)|T| - \sum_{j=2}^{k-1} (\delta^2 + (2 - j)\delta + k + 1 - j)n_j}{\delta^2 + (2 - k)\delta + 1} \leq n_k.$$

Using the Clark et al. [6] notation $a_j = (\delta + 1)(\delta + 1 - j)$, we can write

$$n_k = \left\lceil \frac{kn - (a_1 + k)|T| - \sum_{j=2}^{k-1} (a_j + k)n_j}{a_k + k} \right\rceil.$$

Consider now a recursion similar to that of n_k , but without taking ceilings.

Set $s_2 = \frac{2n - (\delta^2 + d + 2)|T|}{\delta^2 + 1}$ and

$$s_k = \frac{kn - (a_1 + k)|T| - \sum_{j=2}^{k-1} (a_j + k)s_j}{a_k + k}.$$

As the coefficients in the recursions for n_k and s_k are the same as those in Clark et al. [6] (see pages 15–16 in the paper on Clark's webpage), their inequality $n_2 + \dots + n_{\delta+1} \leq s_2 + \dots + s_{\delta+1}$ holds, with the very same proof that they provided.

Following Clark et al. [6], define $u_j = s_j/n$, namely $u_2 = \frac{2}{\delta^2 + 1} - \frac{\delta^2 + \delta + 2}{\delta^2 + 1} \cdot t$ and $u_k = \frac{k - (a_1 + k)t - \sum_{j=2}^{k-1} (a_j + k)u_j}{a_k + k}$. From the displayed formula above, for $k \geq 2$, we have the recurrence

$$k - (a_1 + k)t = \sum_{j=2}^k (a_j + k)u_j.$$

Preparing for an Abel summation, set $S_1 = 0$, $S_k = \sum_{j=2}^k u_j$, $u_j = S_j - S_{j-1}$. Our goal is to compute $S_{\delta+1}$ and use the estimate $|T| + n \cdot S_{\delta+1} \geq \gamma'_6(G)$.

Start with

$$k - (a_1 + k)t = \sum_{j=2}^k (a_j + k)(S_j - S_{j-1}) = S_k(a_k + k) + \sum_{j=2}^{k-1} S_j(\delta + 1).$$

Define now $X_1 = 0$ and $X_k = \sum_{j=2}^k S_j$. Clearly $S_j = X_j - X_{j-1}$,

$$(7) \quad X_2 = \frac{2n - |T|(d^2 + d + 2)}{n(d^2 + 1)}$$

and the displayed formula above turns into

$$k - (a_1 + k)t = (X_k - X_{k-1})(a_k + k) + (\delta + 1)X_{k-1}$$

for $k \geq 3$. Replacing the a -terms with their definition, we obtain the inhomogeneous linear recurrence with polynomial coefficients

$$(8) \quad (\delta^2 + (2 - k)\delta + 1)X_k = \delta(\delta - k + 1)X_{k-1} + k - (\delta^2 + \delta + k)t.$$

In (8) above, setting $k = \delta + 1$, the coefficient of X_{k-1} vanishes, and therefore we obtain

$$X_{\delta+1} = 1 - (\delta + 1)t.$$

We solve the recurrence (8) with initial condition (7). Observe that $-\delta + (1 - t)k$ is a particular solution to (8). The homogeneous recurrence corresponding to (8) is solved by $C \prod_{i=\delta-k+1}^{\delta-2} \left(1 + \frac{\delta+1}{i\delta}\right)^{-1}$ with any C constant. Write

$$X_k = -\delta + (1 - t)k + \frac{\delta - 2(1 - t) + X_2}{\prod_{i=\delta-k+1}^{\delta-2} \left(1 + \frac{\delta+1}{i\delta}\right)}.$$

As this is the sum of a particular solution of (8) and a particular solution ($C = \delta - 2(1 - t) + X_2$) of the homogeneous recurrence corresponding to

(8), this is a solution for (8), which meets the initial condition (7) for $k = 2$. Therefore it is a solution for $2 \leq k \leq \delta$. Finally

$$\begin{aligned} \gamma'_6(G)/n &\leq t + S_{\delta+1} = t + X_{\delta+1} - X_\delta \\ &= t + \left(1 - (\delta + 1)t\right) - \left(-\delta + (1 - t)\delta + \frac{\delta - 2(1 - t) + X_2}{\prod_{i=1}^{\delta-2} \left(1 + \frac{\delta+1}{i\delta}\right)}\right) = A_\delta. \end{aligned}$$

□

To simplify the rest of the proof of Theorem, use the following notation:

$$\begin{aligned} P_d &= \prod_{j=1}^{d-1} \left(1 + \frac{d+1}{jd}\right) \\ Q_d &= 1 - \frac{d-1}{P_d} \\ R_d &= \frac{1}{P_d} \end{aligned}$$

In the next step, we will construct a small dominating set for G . Consider the the following algorithm. In the first step, we find a maximal $T \subseteq V(G)$ with the property that any two vertices in T are of distance at least 3. Note that $N(T)$, the open neighborhood of T is a dominating set. Let $t = |T|/n$, as in Lemma 2.1.1.

If $td < Q_d - tR_d$, then output $N(T)$. Otherwise start a greedy algorithm from T , and output the result.

Observe that $|N(T)| = tdn$. Also, due to Lemma 2.1.1, the greedy algorithm produces a dominating set of size at most $A_d n = (Q_d - tR_d)n$. So the constructed dominating set is the largest, if $td = Q_d - tR_d$. In that case

$$t = \frac{Q_d}{R_d + d} = \frac{P_d - d + 1}{1 + dP_d},$$

and the size of the constructed dominating set is at most

$$tdn = \left(1 - \frac{d^2 - d + 1}{1 + dP_d}\right)n = B_d n.$$

□□

3. DOMINATION IN KNESER GRAPHS

As usual, set $[n] = \{1, 2, \dots, n\}$. The Kneser graph $K(n, k)$ has vertex set $\binom{[n]}{k}$ and vertices $A, B \subseteq [n]$ are adjacent if they are disjoint. The cases $k = 1$ (when $K(n, k)$ is the complete graph on n vertices) and $\frac{n}{2} < k$ (when $K(n, k)$ is the empty graph) are trivial. Domination numbers of Kneser graphs have been studied in the manuscript of Hartman and West [8] and in the Master's Thesis of Gorodezky [7].

Recall that an $S(t, k, v)$ block design is a set of k -subsets (called blocks) of a v -set, where every t -subset of the v -set appears in precisely one block. Surprisingly, the lower bound (3) for the domination number is tight for the graphs $K(7, 3)$ and $K(11, 5)$, as the designs $S(2, 3, 7)$ and $S(4, 5, 11)$ dominate them (see [8], p. 34). So it turns out that the Kneser graphs $K(2k + 1, k)$ form small k are not good examples for regular graphs of large dominating numbers. However we still may have some hope for larger odd graphs, since we do not have corresponding block designs there. It is worth noting that the Lovász Local Lemma does not seem to help at finding regular graphs of large dominating numbers.

Gorodezky [7] showed that for $n > k(k + 1)$ it is true that $\gamma(K(n, k)) = \gamma_t(K(n, k)) = k + 1$. Ivanko and Zelinka [10] showed that for $n \geq 6$, $\text{dom}(K(n, 2)) = \lfloor \frac{1}{3} \binom{n}{2} \rfloor$. We generalize this result below:

Theorem 3.1. *If $n \geq k(k + 1)$ then $\text{dom}(K(n, k)) = \lfloor \frac{1}{k+1} \binom{n}{k} \rfloor$.*

Proof. It is easy to see [7] that a collection of $k + 1$ disjoint k -subsets of $[n]$ gives a (total) dominating set of $K(n, k)$.

Choose $q = \lfloor \frac{1}{k+1} \binom{n}{k} \rfloor$ and let $a_1 = \dots = a_{q-1} = k + 1$. and $a_q = \binom{n}{k} - \sum_{j < q} a_j$. Note that $1 \leq a_q \leq k + 1$ and $a_j = k + 1$ precisely when $j \leq \lfloor \frac{1}{k+1} \binom{n}{k} \rfloor$. Also, $\sum a_j = \binom{n}{k}$. Baranyai's theorem [4] states that there are sets $\mathcal{Y}_1, \dots, \mathcal{Y}_q \subseteq \binom{[n]}{k}$ such that $|\mathcal{Y}_i| = a_i$ and for all i and any $x, y \in [n]$ we have that the number of sets x and y is an element of in \mathcal{Y}_i differs in at most 1. Thus, if $a_i = k + 1$, then \mathcal{Y}_i must be a (total) dominating set. □

REFERENCES

- [1] Noga Alon. Transversal numbers of uniform hypergraphs. *Graphs Combin.*, 6(1):1-4, March 1990.
- [2] Noga Alon and Joel H. Spencer. *The Probabilistic Method*. Wiley, 3 edition, 2008.
- [3] Vladimir Ivanovich Arnautov. Estimation of the exterior stability number of a graph by means of the minimal degrees of the vertices. *Prikl. Mat. i Programirovanie*, 11(126):3-8, 1974.
- [4] Zsolt Baranyai. On the factorization of the complete uniform hypergraph. In *Infinite and finite sets (Colloq., Keszthely, 1973; dedicated to P. Erdős on his 60th birthday)*, Vol. I. North-Holland, Amsterdam, 1975.
- [5] W. Edwin Clark and Larry A. Dunning. Tight upper bounds for the domination numbers of graphs with given order and minimum degree. *Electron. J. Combin.*, 4(1):Research Paper 26, 25 pp. (electronic), 1997.
- [6] W. Edwin Clark, Boris Shekhtman, Stephen Suen, and David C. Fisher. Upper bounds for the domination number of a graph. *Congr. Numer.*, 132:99-123, 1998.
- [7] Igor Gorodezky. Dominating sets in Kneser graphs. Master's thesis, University of Waterloo, 2007. <http://uwspace.uwaterloo.ca/bitstream/10012/3190/1/thesis.pdf>.
- [8] Chris Hartman and Douglas B. West. Covering designs and domination in Kneser graphs. manuscript, 2003.

- [9] Teresa W. Haynes, Stephen T. Hedetniemi, and Peter J. Slater. *Fundamentals of domination in graphs*, volume 208 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker Inc., New York, 1998.
- [10] Jaroslav Ivančo and Bohdan Zelinka. Domination in Kneser graphs. *Math. Bohem.*, 118(2):147–152, 1993.
- [11] Hai Long Liu and Liang Sun. On domination number of 4-regular graphs. *Czechoslovak Math. J.*, 54(129)(4):889–898, 2004.
- [12] László Lovász. On the ratio of optimal and integral fractional covers. *Discrete Math.*, 13:383–390, 1975.
- [13] László Lovász. A kombinatorika minimax tételeiről. *Mat. Lapok*, 26(3-4):209–264, 1975.
- [14] William D. McCuaig and F. Bruce Shepherd. Domination in graphs with minimum degree two. *J. Graph Theory*, 13(6):749–762, 1989.
- [15] Henry Martyn Mulder. *The interval function of a graph*, volume 132 of *Mathematical Centre Tracts*. Mathematisch Centrum, Amsterdam, 1980.
- [16] Oystein Ore. *Theory of graphs*. American Mathematical Society Colloquium Publications, Vol. XXXVIII. American Mathematical Society, Providence, R.I., 1962.
- [17] Charles Payan. Sur le nombre d'absorption d'un graphe simple. *Cahiers Centre Études Recherche Opér.*, 17:307–317, 1975.
- [18] Bruce Reed. Paths, stars and the number three. *Combin. Probab. Comput.*, 5(3):277–295, 1996.
- [19] Hua-Ming Xing, Liang Sun, and Xue-Gang Chen. An upper bound for domination number of 5-regular graphs. *Czechoslovak Math. J.*, 56(131)(3):1049–1061, 2006.

UNIVERSITY OF LOUISVILLE

E-mail address: csaba.biro@louisville.edu

UNIVERSITY OF SOUTH CAROLINA

E-mail address: czabarka@math.sc.edu

UNIVERSITY OF KWAZULU-NATAL

E-mail address: dankelma@ukzn.ac.za

UNIVERSITY OF SOUTH CAROLINA

E-mail address: szekely@math.sc.edu

Applying Integer Programming to Enumerate Equilibrium States of a Multi-Link Inverted Pendulum: A Strange Binomial Coefficient Identity and its Proof

Peter J. Larcombe and Eric J. Fennessey[†]

School of Computing and Mathematics
University of Derby
Kedleston Road, Derby DE22 1GB, U.K.
{P.J.Larcombe@derby.ac.uk}

[†]BAE Systems Integrated System Technologies
Broad Oak, The Airport, Portsmouth PO3 5PQ, U.K.
{Eric.Fennessey@baesystems.com}

Abstract

The task of counting certain (unstable) equilibrium states associated with an idealised 2D inverted pendulum system—some of which describe a particular type of random walk—is re-examined as an optimisation problem. In recovering a known general result, an unusual binomial coefficient identity is generated and requires a non-trivial proof. The complexity of the algorithm leading to state enumeration and identification is, as the number of links increases, discussed in the light of previous work. In addition, a similar type of identity is formulated (and proven) by appealing to some independent analysis by Carlitz from the 1970s, and generalised (computer-based) versions of both identities are included for completeness.